

Varietal Restrictions

Varietal restrictions apply to the grades on this chart. A variety must not be binned above its maximum allowable varietal classification (APH2 > H1/H2 > APW1 > ASW1 > AGP1 > FED1). Only APH2 and H1/H2 varieties may be binned in AUH2 & HPS1.

The varieties Lorikeet, Rosella and Sunsoft 98 must not be binned above AGP1 unless there is an ASWS, ANW1 or ANW2 segregation and they are acceptable varieties at that site.

The varieties Anlace, Barham, Bindawarra, Bowie, Buckley, Corrigan, Datatine, Gazelle, Impala, Orion, QAL2000, QALBis, Snipe, Tatiara, Thornbill, Tincurrin, Wyuna and Yenda must not be binned above AGP1 unless there is a SFE1, SFE2, SGP1 or SGP2 segregation and they are acceptable varieties at that site.

These standards are to be applied on individual truck loads and must not be averaged over a number of loads
Segregations indicated on this chart are only available where announced

Test Code	Binned Grade	APH2	H1	H2	APW1	ASW1	AUH2	AGP1	HPS1	FED1
MOGR	Moisture Maximum (%)	13.5	13.5	13.5	13.5	13.5	13.5	13.5	13.5	13.5
PRGR	Protein (%) Minimum	13.0	13.0	11.5	10.5	No Limit	11.5	No Limit	11.5	No Limit
TWT	Test Weight Minimum (kg/hl)	76	76	76	76	76	71	68	68	62
UNMB	Screenings (Unmillable Material Below the Screen) Maximum (% by weight) Material passing through a 2mm slotted screen after 40 shakes	5.0	5.0	5.0	5.0	5.0	10.0	10.0	25.0	15.0
UNM	Unmillable Material Above the Screen Maximum (% by weight) Material other than wheat kernels remaining on top of a 2.0mm slotted screen after 40 shakes Includes whiteheads (with grains removed), chaff, backbone, Wild Radish pods, Milk Thistle pods, other seedpods less than 5mm in diameter.	0.6	0.6	0.6	0.6	0.6	1.2	1.2	2.6	2.6

Defective Grains – Maximum tolerance % by count per 300 Grain Sample unless otherwise stated (1% = 3 grains), sample to be taken from above the screen.

Defective definitions are to be read in conjunction with the photo in the Visual Recognition Standards Guide which depicts the minimum standard for a grain to be classified as defective.

Note: NIL tolerance applies to the entire load except where stated otherwise.

FALL	Falling Number Minimum (seconds)	350	300	300	300	300	250	200	250	No Limit
SPRO	Sprouted Maximum (% by count) Sprouted kernels are those in which the covering of the germ is split and any further advanced stage of growth to the extent of the germ having grown or shot Note: The NIL tolerance for sprouted grains does not apply if a Falling Number analysis is conducted	NIL	NIL	NIL	NIL	NIL	NIL	NIL	NIL	No Limit
FFUN	Total Stained Maximum (% by count) Grains with a fungal-like discoloration on less than 50% of the entire grain surface (Does not include Storage Moulds) Includes: Blackpoint, Blacktip, Staining due to Moist Plant Material	5% (15 grains)	5% (15 grains)	5% (15 grains)	5% (15 grains)	5% (15 grains)	15% (45 grains)	15% (45 grains)	10% (30 grains)	50% (150 grains)
PFUN	Included in Stained: Pink Stained Maximum (% by count) affected by the fungi (<i>Fusarium</i> , <i>Eppicoccum</i> or <i>Drechslera</i> spp.) Grains that are pink but also contain a white to grey fungal like discoloration over more than 50% of the seed coat surface are to be classified as "White Grain Disorder/Head Scab"	2% (6 grains)	2% (6 grains)	2% (6 grains)	2% (6 grains)	2% (6 grains)	2% (6 grains)	5% (15 grains)	3% (9 grains)	5% (15 grains)
WGDH	Included in Stained: White Grain Disorder/Head Scab Maximum (% by count)	1% (3 grains)	1% (3 grains)	1% (3 grains)	1% (3 grains)	1% (3 grains)	1% (3 grains)	1% (3 grains)	1% (3 grains)	1% (3 grains)
FFAG	Field Fungi Maximum (count per half litre) Grains that are 50% or less discoloured are to be classified as Stained. Grains that are soft and/or emit a mouldy odour are to be classified as Rotted.	10	10	10	10	10	10	10	20	40
GREE	Dry Green or Sappy Maximum (% by count)	1% (3 grains)	1% (3 grains)	1% (3 grains)	1% (3 grains)	1% (3 grains)	2% (6 grains)	5% (15 grains)	2% (6 grains)	No Limit
FTDG	Frost Damaged Maximum (% by count)	1% (3 grains)	1% (3 grains)	1% (3 grains)	1% (3 grains)	1% (3 grains)	2% (6 grains)	10% (30 grains)	2% (6 grains)	No Limit
BURN	Heat Damaged, Bin Burnt and Storage Mould (count per half litre)	1	1	1	1	1	1	1	1	5
SMUT	All Smuts (except Loose Smut) Maximum (entire load) Includes: Ball Smut, Stinking Smut, Bunt, Covered Smut	NIL	NIL	NIL	NIL	NIL	NIL	NIL	NIL	NIL
TAKE	Takeall Affected Maximum (% by count)	1% (3 grains)	1% (3 grains)	1% (3 grains)	1% (3 grains)	1% (3 grains)	2% (6 grains)	1% (3 grains)	2% (6 grains)	No Limit
DAMI	Insect Damaged Maximum (% by count)	1% (3 grains)	1% (3 grains)	1% (3 grains)	1% (3 grains)	1% (3 grains)	1% (3 grains)	2% (6 grains)	2% (6 grains)	4% (12 grains)
ARTD	Over-Dried Damaged Maximum (% by count)	NIL	NIL	NIL	NIL	NIL	NIL	NIL	NIL	No Limit

Foreign Seed Contaminants – Tolerances apply to whole seeds or their equivalent in pieces and refer to the maximum total of all seeds named in each type per half litre.

Except TYPE(1) in which the maximum applies on an individual seed basis per half litre.

WS1	TYPE(1): Colocynth, Double Gees/Spiny Emex/Three Cornered Jack, Jute, Long Head Poppy, Mexican Poppy, New Zealand Spinach, Opium Poppy, Parthenium Weed**, Poppy (Field), Poppy (Horned), Wild Poppy	8*	8*	8*	8*	8*	8*	8*	8*	8*
WS2	TYPE(2): Castor Oil Plant, Coriander, Crow Garlic/Wild Garlic, Darling Pea, Peanut seeds and pods, Ragweed, Rattlepods, Starburr, St. Johns Wort	NIL	NIL	NIL	NIL	NIL	NIL	NIL	NIL	NIL
WS3A	TYPE(3a): Bathurst Burr, Bellvine, Branched Broomrape, Bulls Head/Caltrop/Cats Head, Cape Tulip, Cottonseed, Dodder, Noogoora Burr, Thornapple	2	2	2	2	2	2	2	2	2
WS3B	TYPE(3b): Vetch (Commercial), Vetch (Tare)	4	4	4	4	4	4	4	4	4
WS3C	TYPE(3c): Heliotrope (Blue), Heliotrope (Common), Note included in this Type are tolerances for seeds or pods	2 Pods / 8 Seeds	2 Pods / 8 Seeds	2 Pods / 8 Seeds	2 Pods / 8 Seeds	2 Pods / 8 Seeds	2 Pods / 8 Seeds	2 Pods / 8 Seeds	2 Pods / 8 Seeds	2 Pods / 8 Seeds
WS4	TYPE(4): Bindweed (Field), Cutleaf Mignonette seeds or pods, Darnel (Drake Seed), Hexham Scent/King Island Melilot***, Hoary Cress, Mintweed, Nightshades, Paddy Melon, Skeleton Weed, Variegated Thistle	20	20	20	20	20	20	20	20	20
WS5	TYPE(5): Knapweed (Creeping/Russian), Paterson's Curse/Salvation Jane, Sesbania Pea	40	40	40	40	40	40	40	40	40
WS6	TYPE(6): Colomus Grass, Johnson Grass, Saffron Thistle	10	10	10	10	10	50	50	50	50
WS7A	TYPE(7a): Broad Beans, Chickpeas, Corn (Maize), Cowpea, Faba Beans, Lentils, Lupin, Peas (Field), Safflower, Soybean, Sunflower, And any other seeds or pods greater than 5mm in diameter (excludes milk thistle pods and wild radish pods)	1	1	1	1	1	10	10	10	100
WS7B	TYPE(7b): Barley (2 row), Barley (6 row), Bindweed (Australian), Bindweed (Black), Durum (Unlimited in Feed), Red/Spring Feed Wheats (Unlimited in Feed), Oats (Black or Wild), Oats (Sand), Oats (Common), Rice, Rye (Cereal), Sorghum (Grain), Triticale, Turnip Weed, Type 7b includes any other Foreign Seeds not specified in Types 1 - 7a or elsewhere in Small Foreign Seeds or Unmillable Material Above the Screen.	50	50	50	50	50	150	150	150	400
SFS	Small Foreign Seeds maximum (% by weight) Foreign seeds not specified in TYPE(1)-TYPE(7b) falling below 2mm agtator screen	0.6%	0.6%	0.6%	0.6%	0.6%	1.2%	1.2%	1.2%	1.2%

*Individual Seed Basis ** Parthenium Weed is a NIL tolerance in NSW/VIC/SA ***Hexham Scent is only acceptable if no tainting odour is present

Other Contaminants – Maximum tolerance per half litre unless otherwise stated. - Note: NIL tolerance applies to the entire load except where stated otherwise.

PICK	Pickling Compounds Maximum (entire load)	NIL	NIL	NIL	NIL	NIL	NIL	NIL	NIL	NIL
CHEM	Chemicals Not Approved for Wheat Maximum (entire load)	NIL	NIL	NIL	NIL	NIL	NIL	NIL	NIL	NIL
ERGR	Ryegrass Ergot Maximum (length in cm when pieces are aligned per half litre)	2cm	2cm	2cm	2cm	2cm	2cm	2cm	2cm	2cm
ERGW	Cereal Ergot Maximum (count per half litre)	1	1	1	1	1	1	1	1	1
SGIL	Stored Grain Insects and Pea Weevils: Live maximum (entire load)	NIL	NIL	NIL	NIL	NIL	NIL	NIL	NIL	NIL
INLG	Field Insects – Large, dead or alive Maximum (count per half litre) Includes Desiantha Weevil, Hairy Fungus Beetle, Pea Weevil, Sitona Weevil, Rutherglen bugs, ladybirds, grasshoppers and wood bugs, whole or parts thereof These are insect contaminants of grain that do not cause damage to stored grains	3	3	3	3	3	3	3	3	3
INSM	Field Insects – Small, dead or alive Maximum (count per half litre) Includes all species of aphid, minute mould beetle, mites & stored grain insects (dead only). These are insect contaminants of grain that do not cause damage to stored grains	10	10	10	10	10	10	10	10	10
SNAL	Snails (Live or Dead) Maximum (count per half litre)	1	1	1	1	1	1	10	5	10
EAR	Earcokle Maximum (count per half litre)	10	10	10	10	10	10	15	10	50
LSMT	Loose Smut Maximum (count per half litre)	3	3	3	3	3	3	3	3	3
ESA	Sand Maximum (count per half litre)	20	20	20	20	20	50	50	50	50
ESAP	Earth (5mm maximum diameter) Maximum (count per half litre)	1	1	1	1	1	3	3	1	6
STON	Stones (g per 2.5L) Maximum weight of all Stones retained above a 2.0mm screen per 2.5L	4	4	4	4	4	4	4	4	4
OBJM	Objectionable Material Maximum (entire load) Includes sticks, gravel, odour, tainting agents and other foreign material such as animal excreta, animal meal or carcasses, glass, concrete and metal	NIL	NIL	NIL	NIL	NIL	NIL	NIL	NIL	NIL
ONOM	Other Foreign Material Maximum (% by weight) Includes fine material (dust and minerals), pieces snail shell, sticks less than 1cm long and 0.5cm diameter, other non-vegetative material	0.1%	0.1%	0.1%	0.1%	0.1%	0.1%	0.1%	0.1%	0.2%

This chart is based on Grain Trade Australia (GTA) standards CSG-100, CSG-101, CSG-102, CSG-103, CSG-105, CSG-106, CSG-107, CSG-109 and CSG-150

Not to be assessed at Sample Stand unless otherwise advised

www.graintrade.org.au

GrainFlow Enquiries Toll Free 1800 4GRAIN
www.grainflow.com.au